Cluster of Laslett

| Patient information | | |
|--|------------------------|---|
| Name: | | |
| Age: | | |
| Gender: | | |
| Medical history: | | |
| | | |
| | | |
| | | |
| Primary complaint: | | |
| | | |
| | | |
| | | |
| Instructions for clinicians | | |
| Apply pressure gradually and ensure the patient's safety. | | |
| Accurately note the test outcomes to confirm the diagnosis. | | |
| Stop testing once 2/4 of tests are positive or proceed if necessary. | | |
| Consider referral for needed. | r fluoroscopy-guided r | nerve block if pain persists or further confirmation is |
| Provocative tests | | |
| Distraction test | | |
| Outcome: | Positive | Negative |
| Thigh thrust test | | |
| Outcome: | Positive | Negative |
| Compression test | | |
| Outcome: | Positive | Negative |
| Sacral thrust test | | |
| Outcome: | Positive | Negative |
| Gaenslen's test (optional, supplementary) | | |
| Outcome: | Positive | Negative |

| Evaluation |
|--|
| Test positives: |
| Distraction |
| Thigh thrust |
| Compression |
| Sacral thrust |
| Gaenslen's |
| Diagnosis (Cluster of Laslett) |
| Positive sacroiliac joint dysfunction: |
| Further testing required: |
| Notes |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| Clinician information |
| Name: |
| Credentials: |
| Signature: |
| Date: |