

# Impetigo Diagnosis Handout

Impetigo is a common bacterial skin infection that most commonly affects children. It can be easily spread through close physical contact or by sharing items such as towels, clothing, or toys. Impetigo typically appears as red sores on the face, especially around the mouth and nose area. In this handout, we will explain how impetigo is diagnosed and provide some tips for prevention.

The two types of impetigo are nonbullous impetigo (i.e., impetigo contagiosa) and bullous impetigo.

## Diagnosing impetigo

The diagnosis of nonbullous and bullous impetigo is nearly always clinical. Differential diagnosis includes many other blistering and rash disorders. Skin swabs cannot differentiate between bacterial infection and colonization.

In patients in whom first-line therapy fails, culture of the pus or bullous fluid, not the intact skin, may be helpful for pathogen identification and antimicrobial susceptibilities.

Although serologic testing for streptococcal antibodies is helpful in the diagnosis of acute poststreptococcal glomerulonephritis, it does not aid in the diagnosis of impetigo.

## Differential diagnosis

Impetigo should be differentiated from other common skin conditions such as eczema, herpes simplex virus infection, and candidiasis. In some cases, a bacterial culture or skin biopsy may be necessary to confirm the diagnosis.

Below are some details about the differential diagnosis of impetigo:

### Bullous impetigo

Diagnosis	Distinguishing features
<b>Bullous erythema multiforme</b>	Vesicles or bullae arise from a portion of red plaques, 1 to 5 cm in diameter, on the extensor surfaces of extremities.
<b>Bullous lupus erythematosus</b>	Widespread vesiculobullous eruption that may be pruritic; tends to favor the upper part of the trunk and proximal upper extremities.
<b>Bullous pemphigoid</b>	Vesicles and bullae appear rapidly on widespread pruritic, urticarial plaques.
<b>Herpes simplex virus</b>	Grouped vesicles on an erythematous base that rupture to become erosions covered by crusts, usually on the lips and skin; may have prodromal symptoms.
<b>Insect bites</b>	Bullae seen with pruritic papules grouped in areas where bites occur.
<b>Pemphigus vulgaris</b>	Nonpruritic bullae, varying in size from 1 to several centimeters, appear gradually and become generalized; erosions last for weeks before healing with hyperpigmentation, but no scarring occurs.
<b>Stevens-Johnson syndrome</b>	Vesiculobullous disease of the skin, mouth, eyes, and genitalia; ulcerative stomatitis with hemorrhagic crusting is the most characteristic feature.
<b>Thermal burns</b>	History of burn with blistering in second-degree burns.

Diagnosis	Distinguishing features
<b>Toxic epidermal necrolysis</b>	Stevens-Johnson–like mucous membrane disease followed by diffuse generalized detachment of the epidermis.
<b>Varicella</b>	Thin-walled vesicles on an erythematous base that start on the trunk and spread to the face and extremities; vesicles break and crusts form; lesions of different stages are present at the same time in a given body area as new crops develop.

## Nonbullous impetigo

Diagnosis	Distinguishing features
<b>Atopic dermatitis</b>	Chronic or relapsing pruritic lesions and abnormally dry skin; flexural lichenification is common in adults; facial and extensor involvement is common in children
<b>Candidiasis</b>	Erythematous papules or red, moist plaques; usually confined to mucous membranes or intertriginous areas
<b>Contact dermatitis</b>	Pruritic areas with weeping on sensitized skin that comes in contact with haptens (e.g., poison ivy)
<b>Dermatophytosis</b>	Lesions may be scaly and red with slightly raised “active border” or classic ringworm; or may be vesicular, especially on feet
<b>Discoid lupus erythematosus</b>	Well-defined plaques with adherent scale that penetrates into hair follicles; peeled scales have “carpet tack” appearance
<b>Ecthyma</b>	Crusted lesions that cover an ulceration rather than an erosion; may persist for weeks and may heal with scarring as the infection extends to the dermis
<b>Herpes simplex virus</b>	Vesicles on an erythematous base that rupture to become erosions covered by crusts, usually on the lips and skin
<b>Insect bites</b>	Papules usually seen at site of bite, which may be painful; may have associated urticaria
<b>Pemphigus foliaceus</b>	Serum and crusts with occasional vesicles, usually starting on the face in a butterfly distribution or on the scalp, chest, and upper back as areas of erythema, scaling, crusting, or occasional bullae
<b>Scabies</b>	Lesions consist of burrows and small, discrete vesicles, often in finger webs; nocturnal pruritus is characteristic
<b>Sweet’s syndrome</b>	Abrupt onset of tender or painful plaques or nodules with occasional vesicles or pustules
<b>Varicella</b>	Thin-walled vesicles on an erythematous base that start on trunk and spread to face and extremities; vesicles break and crusts form; lesions of different stages are present at the same time in a given body area as new crops develop

Cole, C., & Gazewood, J. (2007). Diagnosis and treatment of impetigo. *American Family Physician*, 75(6), 859–864. <https://pubmed.ncbi.nlm.nih.gov/17390597/>

Hartman-Adams, H., Banvard, C., & Juckett, G. (2014). Impetigo: Diagnosis and treatment. *American Family Physician*, 90(4), 229–235. <https://www.aafp.org/pubs/afp/issues/2014/0815/p229.html>