## **Quadriceps Active Test**

Patient information
Name:
Date of birth:
Gender:
Date of test:
Height:
Weight:
Test overview
The Quadriceps Active Test is a clinical examination used to detect a tear of the knee's posterior cruciate ligament (PCL). This test is beneficial for patients who present with knee pain, instability, or a history of trauma to the knee. The test helps determine the integrity of the PCL by assessing the anterior translation of the tibia when the quadriceps muscle is contracted.
Instructions for the patient
Wear comfortable clothing that allows clear visibility of the knee.
Lie on your back with your legs fully extended.
Follow the clinician's instructions carefully during the test.
Test procedure
1. The patient lies supine (on their back) on the examination table with both legs extended.
2. The clinician bends the affected knee to 90 degrees while the foot remains flat on the table.
3. The clinician stabilizes the patient's foot and asks the patient to relax their quadriceps muscle.
4. The clinician instructs the patient to contract the quadriceps muscle while applying resistance to the lower leg.
5. The clinician observes the anterior translation of the tibia. An excessive forward movement of the tibia indicates a positive Quadriceps Active Test, suggesting a PCL injury.
Measurements
Affected knee
Anterior translation of the tibia:
Knee flexion angle:
Quadriceps muscle contraction:
Present Absent

Unaffected knee
Anterior translation of the tibia:
Knee flexion angle:
Pain level during test (0-10):
Quadriceps muscle contraction:
Present Absent
Interpretation
Positive
Indication: Excessive forward movement of the tibia on the affected side.
Possible diagnosis: Posterior Cruciate Ligament (PCL) tear.
• Suggests a PCL injury, which may require further imaging (MRI) and orthopedic consultation.
<ul> <li>Indicates a potential need for physical therapy focusing on knee stabilization and strengthening.</li> </ul>
Negative
Indication: No excessive forward movement of the tibia.
Possible diagnosis: Unlikely PCL tear; consider other potential knee issues.
Healthcare practitioner's notes
Additional observations:
Additional observations:
Additional observations:
Additional observations:  Patient feedback:
Patient feedback:
Patient feedback:
Patient feedback:  Recommendations and follow-up plan:
Patient feedback:  Recommendations and follow-up plan:  Healthcare practitioner's information
Patient feedback:  Recommendations and follow-up plan:  Healthcare practitioner's information  Name: